

A new approach to q -zeta function

Taekyun Kim

Institute of Science Education,
Kongju National University, Kongju 314-701, Korea
e-mail: tkim@kongju.ac.kr

Abstract We construct the new q -extension of Bernoulli numbers and polynomials in this paper. Finally we consider the q -zeta functions which interpolate the new q -extension of Bernoulli numbers and polynomials.

2000 Mathematics Subject Classification - 11B68, 11S40, 11S80

Key words- Bernoulli numbers and polynomials, zeta functions

1. Introduction

Throughout this paper $\mathbb{Z}, \mathbb{Z}_p, \mathbb{Q}_p$ and $\mathbb{C}_p$ will be denoted by the ring of rational integers, the ring of p -adic integers, the field of p -adic rational numbers and the completion of algebraic closure of $\mathbb{Q}_p$, respectively, cf. [7, 8, 9, 10]. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$. When one talks of q -extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$, or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}_p$, then we normally assume $|q - 1|_p < p^{-\frac{1}{p-1}}$, so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$. If $q \in \mathbb{C}$, then we normally assume that $|q| < 1$. For $f \in UD(\mathbb{Z}_p, \mathbb{C}_p) = \{f|f : \mathbb{Z}_p \rightarrow \mathbb{C}_p \text{ is uniformly differentiable function}\}$, the p -adic q -integral (or q -Volkenborn integration) was defined as

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \quad (1)$$

where $[x]_q = \frac{1 - q^x}{1 - q}$, cf. [1, 2, 3, 4, 11]. Thus we note that

$$I_1(f) = \lim_{q \rightarrow 1} I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_1(x) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{0 \leq x < p^N} f(x), \text{ cf. [4, 11].} \quad (2)$$

By (2), we easily see that

$$I_1(f_1) = I_1(f) + f'(0), \text{ cf. [5, 6, 7],} \quad (3)$$

where $f_1(x) = f(x+1)$, $f'(0) = \frac{d}{dx}f(x)|_{x=0}$.

In [8], the q -Bernoulli polynomials are defined by

$$\beta_n^{(h)}(x, q) = \int_{\mathbb{Z}_p} [x + x_1]_q^n q^{x_1(h-1)} d\mu_q(x_1), \text{ for } h \in \mathbb{Z}. \quad (4)$$

In this paper we consider the new q -extension of Bernoulli numbers and polynomials. The main purpose of this paper is to construct the new q -extension of zeta function and L -function which interpolate the above new q -extension of Bernoulli numbers at negative integer.

2. On the New q -Extensions of Bernoulli numbers and polynomials

In (3), if we take $f(x) = q^{hx}e^{xt}$, then we have

$$\int_{\mathbb{Z}_p} q^{hx} e^{xt} d\mu_1(x) = \frac{h \log q + t}{q^h e^t - 1}. \quad (5)$$

for $|t| \leq p^{-\frac{1}{p-1}}$, $h \in \mathbb{Z}$.

Let us define the (h, q) -extension of Bernoulli numbers and polynomials as follows:

$$\begin{aligned} F_q^{(h)}(t) &= \frac{h \log q + t}{q^h e^t - 1} = \sum_{n=0}^{\infty} B_{n,q}^{(h)} \frac{t^n}{n!}, \\ F_q^{(h)}(t, x) &= \frac{h \log q + t}{q^h e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q}^{(h)}(x) \frac{t^n}{n!}. \end{aligned} \quad (6)$$

Note that $B_{n,q}^{(h)}(0) = B_{n,q}^{(h)}$, $\lim_{q \rightarrow 1} B_{n,q}^{(h)} = B_n$, where B_n are the n -th Bernoulli numbers. By (5) and (6), we obtain the following Witt's formula.

Theorem 1. For $h \in \mathbb{Z}$, $q \in \mathbb{C}_p$ with $|1 - q|_p \leq p^{-\frac{1}{p-1}}$, we have

$$\begin{aligned} \int_{\mathbb{Z}_p} q^{hx} x^n d\mu_1(x) &= B_{n,q}^{(h)}, \\ \int_{\mathbb{Z}_p} q^{hy} (x+y)^n d\mu_1(y) &= B_{n,q}^{(h)}(x). \end{aligned} \quad (7)$$

By above theorem, we easily see that

$$B_{n,q}^{(h)}(x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} B_{k,q}^{(h)}. \quad (8)$$

Let d be any fixed positive integer with $(p, d) = 1$. Then we set

$$\begin{aligned} X &= X_d = \varprojlim_N (\mathbb{Z}/dp^N\mathbb{Z}), X_1 = \mathbb{Z}_p, \\ X^* &= \bigcup_{0 < a < dp} a + dp\mathbb{Z}_p, \\ a + dp^N\mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ with $0 \leq a < dp^N$. Note that

$$\int_{\mathbb{Z}_p} f(x) d\mu_1(x) = \int_X f(x) d\mu_1(x),$$

for $f \in UD(\mathbb{Z}_p, \mathbb{C}_p)$, cf. [1, 2, 3, 4, 10].

In Eq. (7), it is easy to see that

$$\begin{aligned} B_{k,q}^{(h)}(x) &= \int_X (x+t)^k q^{ht} d\mu_1(t) = \lim_{l \rightarrow \infty} \frac{1}{mp^l} \sum_{n=0}^{mp^l-1} q^{hn} (x+n)^k \\ &= \frac{1}{m} \lim_{l \rightarrow \infty} \frac{1}{p^l} \sum_{i=0}^{m-1} \sum_{n=0}^{p^l-1} q^{h(i+mn)} (x+i+mn)^k = m^{k-1} \sum_{i=0}^{m-1} q^{hi} B_{k,q^m}^{(h)} \left(\frac{x+i}{m} \right). \end{aligned}$$

Therefore we have the below theorem.

Theorem 2. For any positive integer m , we have

$$B_{k,q}^{(h)}(x) = m^{k-1} \sum_{i=0}^{m-1} q^{hi} B_{k,q^m}^{(h)} \left(\frac{x+i}{m} \right), \text{ for } k \geq 0.$$

Let χ be the Dirichlet character with conductor $d \in \mathbb{Z}_+$. Then we define the (h, q) -extension of generalized Bernoulli numbers attached to χ . For $n \geq 0$, define

$$B_{n,q,\chi}^{(h)} = \int_X \chi(x) q^{hx} x^n d\mu_1(x). \quad (9)$$

By (9), we easily see that

$$\begin{aligned} B_{n,q,\chi}^{(h)} &= \lim_{l \rightarrow \infty} \frac{1}{dp^l} \sum_{x=0}^{dp^l-1} \chi(x) q^{hx} x^n = \frac{1}{d} \lim_{l \rightarrow \infty} \frac{1}{p^l} \sum_{i=0}^{d-1} \sum_{x=0}^{p^l-1} \chi(i+dx) q^{h(i+dx)} (i+dx)^n \\ &= \frac{1}{d} \sum_{i=0}^{d-1} \chi(i) q^{hi} \int_{\mathbb{Z}_p} q^{hdx} (i+dx)^n d\mu_1(x) = d^{n-1} \sum_{i=0}^{d-1} \chi(i) q^{hi} B_{n,q^d}^{(h)} \left(\frac{i}{d} \right). \end{aligned}$$

Therefore we obtain the below lemma.

Lemma 3. For $d \in \mathbb{Z}_+$, we have

$$B_{k,q,\chi}^{(h)} = d^{k-1} \sum_{i=0}^{d-1} \chi(i) q^{hi} B_{k,q^d}^{(h)} \left(\frac{i}{d} \right), \text{ for } n \geq 0. \quad (10)$$

By induction in Eq.(3), we easily see that

$$I_1(f_b) = I_1(f) + \sum_{i=0}^{b-1} f'(i), \text{ where } f_b(x) = f(x+b), b \in \mathbb{Z}_+. \quad (11)$$

In Eq.(11), if we take $f(x) = q^{hx} e^{tx} \chi(x)$, then we have

$$I_1(e^{tx} q^{hx} \chi(x)) = \frac{\sum_{i=0}^{d-1} (te^{it} \chi(i) q^{hi} + e^{ti} (h \log q) q^{hi} \chi(i))}{q^{hd} e^{dt} - 1}. \quad (12)$$

By (12) and (9), we can give the generation function of $B_{n,q,\chi}^{(h)}$ as follows:

$$F_{q,\chi}^{(h)}(t) = \frac{\sum_{i=0}^{d-1} (te^{it} \chi(i) q^{hi} + e^{ti} (h \log q) q^{hi} \chi(i))}{q^{hd} e^{dt} - 1} = \sum_{n=0}^{\infty} B_{n,q,\chi}^{(h)} \frac{t^n}{n!}. \quad (13)$$

3. The analogue of zeta function

In this section we assume that $q \in \mathbb{C}$ with $|q| < 1$. Let $\Gamma(s)$ be the gamma function. By (6), we can readily see that

$$\begin{aligned} \frac{1}{\Gamma(s)} \int_0^\infty t^{s-2} e^{-t} F_q^{(h)}(-t) dt &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-2} e^{-t} \left\{ \frac{-t}{q^h e^{-t} - 1} + \frac{h \log q}{q^h e^{-t} - 1} \right\} dt \\ &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-t} \frac{1}{1 - q^h e^{-t}} dt - \frac{h \log q}{\Gamma(s)} \int_0^\infty \frac{t^{s-2} e^{-t}}{1 - q^h e^{-t}} dt \\ &= \sum_{n=0}^{\infty} q^{nh} \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-(n+1)t} dt - h \log q \sum_{n=0}^{\infty} q^{nh} \int_0^\infty t^{s-1} e^{-(n+1)t} dt \\ &= \sum_{n=1}^{\infty} \frac{q^{(n-1)h}}{n^s} - \frac{\log q^h}{s-1} \sum_{n=1}^{\infty} \frac{q^{(n-1)h}}{n^{s-1}}. \end{aligned} \quad (14)$$

Using (14), we define the new q -extensions of zeta functions as follows:

Definition 4. For $s \in \mathbb{C}, x \in \mathbb{R}^+$, we define

$$\zeta_q^{(h)}(s) = \sum_{n=1}^{\infty} \frac{q^{(n-1)h}}{n^s} - \frac{h \log q}{s-1} \sum_{n=1}^{\infty} \frac{q^{(n-1)h}}{n^{s-1}}, \quad (14-a)$$

$$\zeta_q^{(h)}(s, x) = \sum_{n=0}^{\infty} \frac{q^{nh}}{(n+x)^s} - \frac{h \log q}{s-1} \sum_{n=0}^{\infty} \frac{q^{nh}}{(n+x)^{s-1}}. \quad (14-b)$$

Remark. By (14-a) and (14-b), we easily see that $\zeta_q^{(h)}(s) = \zeta_q^{(h)}(s, 1)$. Also, we note that $\zeta_q^{(h)}(s)$ is analytic continuation for $R(s) > 1$.

Using Mellin transforms in Eq.(6), we obtain

$$\frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2} F_q^{(h)}(-t, x) dt = \zeta_q^{(h)}(s, x). \quad (15)$$

By (15) and (6), we readily see that

$$\zeta_q^{(h)}(s, x) = \sum_{n=0}^{\infty} \frac{(-1)^n B_{n,q}^{(h)}(x)}{n!} \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2+n} dt.$$

Therefore we obtain the following:

Theorem 5. For $n \in \mathbb{N}$, we have

$$\zeta_q^{(h)}(1-n, x) = -\frac{B_{n,q}^{(h)}(x)}{n}$$

By Mellin transforms and (13), we note that

$$\frac{1}{\Gamma(s)} \int_0^{\infty} F_{q,\chi}^{(h)}(-t) t^{s-2} dt = \sum_{n=1}^{\infty} \frac{q^{nh} \chi(n)}{n^s} - \frac{h \log q}{s-1} \sum_{n=1}^{\infty} \frac{q^{nh} \chi(n)}{n^{s-1}}. \quad (16)$$

Thus we can define the new q -extension of Dirichlet L -function as follows:

Definition 6. For $s \in \mathbb{C}$, we define

$$L_q^{(h)}(s, \chi) = \sum_{n=1}^{\infty} \frac{q^{nh} \chi(n)}{n^s} - \frac{h \log q}{s-1} \sum_{n=1}^{\infty} \frac{q^{nh} \chi(n)}{n^{s-1}}.$$

By (16) and (13), we have the following

Theorem 7. For $n \in \mathbb{N}$, we obtain

$$L_q^{(h)}(1-n, \chi) = -\frac{B_{n,q,\chi}^{(h)}}{n}.$$

References

- [1] T. Kim, Analytic continuation of multiple q -zeta functions and their values at negative integers, *Russian J. Math. Phys.*, **11**(2004), 71-76.
- [2] T. KIM, Non-Archimedean q -integrals associated with multiple Changhee q -Bernoulli polynomials , *Russ. J. Math. Phys.*, **10** (2003), 91-98.
- [3] T. Kim, On Euler-Barnes multiple zeta functions, *Russian J. Math. Phys.*, **10**(2003), 261-267.
- [4] T. Kim, q -Volkenborn integration , *Russ. J. Math. Phys.*, **9**(2002), 288-299.
- [5] T. Kim, An invariant p -adic Integral associated with Daehee numbers, *Integral Transforms and Special Functions* , **13** (2002), 65-69.
- [6] T. Kim, p -adic q -integrals associated with the Changhee-Barne's q -Bernoulli polynomials , *Integral Transforms and Special Functions*, **15**(5)(2004), 415-420.
- [7] T. Kim, q -Riemann zeta functions , *Int. J. Math. Math. Sci.*, **2004**(12) (2004), 599-605.
- [8] T. Kim, S. H. Rim, Generalized Carlitz's q -Bernoulli numbers in the p -adic number field , *Advan. Stud. Contemp. Math.*, **2**(2000), 9-19.
- [9] T. Kim, A note on multiple zeta function , *JP. J. Algebra, Number Theory and Applications*, **3**(3)(2003), 471-476.
- [10] T. Kim et al, Introduction to Non-Archimedean Analysis, Kyo Woo Sa (Korea), 2004. (<http://www.kyowoo.co.kr>).
- [11] T. Kim, A note on Dirichlet L -series, *Proc. Jangjeon Math. Soc.*, **6**(2003), 161-166.